



The Biological and Medical Properties of *Myrtus communis* Linn - Subject Review

1. Sunbul Jalal Shakour

Received 20th Oct 2023,
Accepted 15th Nov 2023,
Online 16th Dec 2023

¹ Collage of Nursing, University of
Kirkuk, Iraq.
sunalsaffar9@uokirkuk.edu.iq

Abstract: The plant called *Myrtus communis* that belong to family of Myrtaceae is known as a significant plant being used in System of Unani of Medicine since ancient Greece time. Typically, *M. communis* is planted for its attractive foliage, the flowers and also the berries. Its berries, foliage and the essential oil are often utilized to various diseases as the ulcer of stomach, diarrhea condition, and the dysentery condition with vomiting, the rheumatism disorders, haemorrhages problems and leucorrhoea. The foliage and the berries are utilized in flavoring. Ripe fruits have historically been used as food additives because of the high vitamin content of this plant. The chemical makeup of *M. communis* includes tannins, phenols, flavonoids, coumarins, essential oils, different types of fiber, sugars, vitamin C, malic acid, and antioxidants. This paper was showed the pharmacological qualities, chemical composition, and medical applications of the plant.

Key words: *M. communis*, Biological activities, Medical activities, Phenols.

Introduction

Myrtus communis, commonly known as "Murt" or "Murd" in traditional texts, is one of the most frequently cited *Myrtus* species [1]. *M. communis*, also referred to as common myrtle, is a Mediterranean shrub that is indigenous to that area. The plant has branches that form a thickly covered entire head and grows to a height of 2.4 to 3 meters [2]. Asif et al. [3] describe the fruits as tiny and black, with small green leaves. The evergreen leaves measure 2 to 5 cm in length. Because of its astringent qualities, it has a bitter flavor [4-6]. The flowers are extremely fragrant, star-shaped, and white

or pinkish [7]. The round, fruit is highly seeded. Flowers are pollinated by insects, and birds who eat berries distribute seeds. (Figure 1) [8]. Long used as an ethnobotanical remedy, myrtle is also used to cure obesity, hypercholesterolemia, and diabetes [11–12], as well as to treat skin conditions, colds and coughs, digestive issues, and diabetes [9–10]. In perfumery, Essential oils derived from shoots, leaves, and occasionally flowers and fruits are frequently used. The berries are also used to make bitters and well-known liquors. Myrtle has been extensively researched due to its widespread use in folk medicine, particularly in terms of EO content [13–19]. Contributions to the literature on biological activity include research on antioxidant and antibacterial properties [20–27], as well as putative anti-inflammatory and antitumoral actions [21–22]. The leaf EO has an antiparasitic action against *Leishmania tropica* strains, as Mahmoudvand et al. [28] have found. Recent studies have found that the fruit's high phospholipid, polyunsaturated fatty acid, and phenolic content accounts for its effectiveness in treating ulcerative colitis, esophagitis, diarrhea, and digestive illnesses [29]. However, the chemical makeup of plant species' extracts, which varies greatly depending on a variety of variables including growing region, season, and cultivar [30], as well as the surrounding conditions, plant phenological phases, and genotypes [31], strongly influences the biological activities of plant species. The same species, which is represented by a diversity of morphs and genotypes, is capable of producing a range of extracts with different compositions and therapeutic effects depending on these characteristics.

Kingdom Plantae – Plants

Division Magnoliophyta

Class Magnoliopsida – Dicotyledons

Subclass Rosidae

Order Myrtales

Family Myrtaceae

Genus *Myrtus*

Species *Myrtus communis* L.

Phytochemical constituents

The plant contains fibers, carbohydrates, antioxidants, and a variety of biologically active chemicals [32–33]. The main phytochemicals present in fruits include anthocyanins, flavonoids, and phenolic compounds. Fixed oil comprises about 12–15% of the fat in seeds and is composed of glycerides comprising oleic acid, the myristic acid, palmitic, and lauric acids [34]. According to fatty acid assays [35], the myrtle fruit contains 14 fatty acids, with oleic acid being the most frequent, followed by palmitic acid and stearic acid. The essential oil of *M. communis*, known as myrtle oil, is obtained by steam distilling the plant's leaves, the fruits, and flowers. Its color is golden or greenish yellow, and it smells distinctly reviving. The yield and quality of oil are determined by the region of production, the harvest season, and the length of distillation. variable sections of the plant have variable oil yields (w/w). The yields of the hydro-distilled oils were as follows: 0.4–0.5 for leaves, 0.4 for flowers, 0.5 for unripe fruits, and 0.02% for ripe fruits. Terpenes and terpene alcohols make for over 80% of the volatile

substances present in essential oils [36]. There are different concentrations of the five terpenoid compounds in leaf oil, fruits, and flowers [37].



Figure (1): branches and fruit of *M. communis*

Pharmacological activities

Anti-bacterial effect

The therapeutic benefits of *Berberis vulgaris* 5% (in metronidazole basis) and *Myrtus communis* L 2% (in metronidazole base) vaginal gels on bacterial vaginosis were evaluated and compared to metronidazole vaginal gel 0.75%. The study's findings suggest that combining *Myrtus communis* and metronidazole improves the efficacy of bacterial vaginosis treatment [38]. Pinene, 1, 8 cineole, and limonene are the most significant contributors to the antimicrobial properties of *M. communis* essential oil, according to research on the antimicrobial efficacy of the samples against three microbial strains. Our findings are supported by other research that has discovered that plant essential oils rich in these compounds have strong antibacterial action [39]. Sodium hypochlorite (NaOCl) and chlorhexidine were compared to this plant's antibacterial activity against *E. faecalis*, *S. aureus*, and *C. albicans*. With a minimum inhibitory concentration in the range of 0.032-32 g/mL, *M. Communis* essential oil was a powerful antibacterial agent against persistent endodontic germs [40].

Antifungal activity

At a dosage of 1600 ppm in vitro, essential oils from the base plant had an inhibitory action against the *R. Solani* fungus of up to 60%. In a study using the broth microdilution method, Investigations were made into the essential oils from the base plant's ability to suppress the *Aspergillus* fungus. The researchers reported an effective impact against *Aspergillus* fungi [41]. *Amaranthus retroflexus*, *Cirsium arvense*, *Lactuca serriola* and *Rumex crispus* seeds were examined for the oils' herbicidal effects on seed germination and seedling growth. The oils prevented plant seed germination and seedling growth

completely or partially. According to the findings of this study, *M. communis* essential oils could be employed as natural fungicides and herbicides [42].

Antioxidant Activity

Various studies on the antioxidant properties of *Myrtus communis* extracts have also been conducted. According to Serce et al., [43], methanolic extracts of eight Turkish myrtle fruit accessions displayed variable degrees of capacity to scavenge DPPH free radicals and inhibit linoleic acid oxidation as measured by the β -carotene-bleaching test. The IC₅₀ values found in the DPPH assay varied from 2.34 g/mL, which was not statistically different from the reference α -tocopherol, to 8.24 g/mL. The inhibition percentages for inhibiting linoleic acid oxidation were consistently above 80% but less than the 96.31% registered for the reference substance (α -tocopherol). Polat et al. [44] investigated how extracting solvents affected the total phenolic content, antioxidant activity, and antiradical action of myrtle extracts collected in different locations of Turkey. [45] examined the antioxidant activity of extracts from white and dark blue Tunisian berries. Tuberoso et al. [46] conducted a first-ever evaluation of the antioxidant activity of myrtle berry extracts made with different polarity solvents. The researchers evaluated the extracts' capacity to neutralize DPPH free radicals and protect biological molecules using cholesterol and LDL (low density lipoproteins) oxidation assays.

Cardiovascular activity

The aqueous leaf extract produced a bad inotropic impact on guinea pigs that wasn't stopped by atropine. The entire extract had a concentration-dependent depressive effect in anesthetized rabbits that was inhibited by theophylline but not attenuated by propranolol, cimetidine, or atropine. These results support the discovery by Al-Zohry et al. of an adenosine-like molecule in this extract [47].

Hepatoprotective activity

It was investigated whether the aqueous extract of *M. Communis* leaves might shield albino rat livers from the toxicity paracetamol caused. A series of biochemical measurements, including alanine transaminase, aspartate transaminase and phosphatase, total protein, and bilirubin level in the blood (both total and direct), were used to determine the extent of liver protection provided by the aqueous extract of *M. Communis* leaves. The liver's protective effects were seen at doses of 200 mg/kg and 400 mg/kg of body weight because the extract significantly decreased the amount of enzymes at these concentrations [48].

The effect on the peptic ulcers

H. pylori infection plays significant role in pathogenesis of gastritis, such as disease called peptic ulcer and the stomach cancer [49-50]. Different studies and researches have been done to determine the protective impact of *M. communis* versus gastric ulcers disease [51]. Rats were given the aqueous extract prior to exposure to ulcerogenic drugs in order to determine the protective effect of myrtle on stomach ulcer versus pyloric ligation-induced models. Aqueous extract treatment considerably lowers the ulcer index. In comparison to omeprazole, the aqueous extract's action is more pronounced at low doses. When

indomethacin, which produces ulcers, was administered to rats' stomachs, histopathological analysis revealed significant anti-ulcer effects at both dose levels [52].

Antidiabetic effects

In mice with diabetes brought on by streptozotocin, *M. communis* leaves were found to have an anti-diabetic effect. Recently, it was discovered that the ace plant's phenolic chemicals were in charge of this activity. *M. Communis* leaves were used to extract phenols, which had an anti-diabetic activity [53]. The usefulness of *M. Communis* in treating diabetic rats did not show any promising results. The outcomes also revealed that ace plant phenols had a considerable anti-diabetic influence when supplied 800 mg/kg body weight and a mild reaction as an anti-diabetic at a dose of 400 mg/kg in rats without diabetes. These effects were demonstrated by regeneration alterations in the cells of the pancreatic island. According to the study, the absence of tissue abnormalities in control mice without diabetes' liver and kidneys revealed that phenolic compounds alone have no adverse impact on physiological tissues [53].

Insecticidal activity

Essential oils from *M. Communis* leaves and flowers were found to have fatal and poisonous effects on mosquito larvae *Culex pipiens* when used as insecticides [54]. Essential oil were exhibit insecticide impact versus 3 stored insects, adults of Mediterranean flour moth, Indian meal moth and *A. getectus* Say bean [55]. Essential oils in the ace plant have been shown to be effective as an insecticide against *P. humanis capitis*. The efficacy as an insecticide may be due to Lineol, α -pinene and the linalool [56].

Reference

1. Mahboubi, M. (2017). Effectiveness of *Myrtus communis* in the treatment of hemorrhoids. *Journal of Integrative Medicine*, 15(5), 351–358.
2. Sumbul, S., Ahmad, M. A., Asif, M., & Akhtar, M. (2011). *Myrtus communis* Linn.—A review. *Indian Journal of Natural Products and Resources*, 2(4), 395–402.
3. Asif, H., Akram, M., Uddin, S., Hasan, Z. U., Sami, A., Iqbal, A., Tauseef, U., & Bari, A. (2011). *Myrtus communis* Linn. (pharmacological activity). *Journal of Medicinal Plants Research*, 5(26), 6257–6259.
4. AL-dalwi L M S, Alaa H. A. Q. (2022). Effect of shading ratio, benzyl adenine and organic fertilizer (Horti boost 10) of Myrtle Plants (*Myrtus communis* L). *IOP Conf. Series: Earth and Environmental Science*. 1158: 1-10.
5. AL-Saiedy A..N. K. and AL-Nuaimy M..I.A. (2019). Effect of adding types of red pepper powder (Bio active subctance) and fibrolytic enzymes mixture in improvement of the performance and quality egg traits of the layer (ISA BROWN), *Kirkuk University Journal For Agricultural Sciences KUJAS*. 2018: Special Issue:203-215.
6. Sadeq, M., & AL-Neemi, M. (2023). The effect of adding local myrtle leaves powder and industrial antioxidant Butylated Hydroxy Toluene to laying hens feeding on some

- biochemical characteristics and special enzymes against oxidative stress in laying hens blood. Kirkuk University Journal For Agricultural Sciences, 14(2), 91-98.
7. Charles, D. J. (2012). Antioxidant properties of spices, herbs and other sources. Springer Science & Business Media.
 8. Satyavati, G., Raina, M., & Sharma, M. (1987). Medicinal plants of India (Vol. 2). Indian Council of Medical Research.
 9. Loi, M.C.; Maxia, L.; Maxia, A. Ethnobotanical comparison between the villages of Escolca and Lotzorai (Sardinia, Italy). J. Herbs Spices Med. Plants 2005, 11, 67–84.
 10. Maxia, A.; Lancioni, M.C.; Balia, A.N.; Alborghetti, R.; Pieroni, A.; Loi, M.C. Medical ethnobotany of the Tabarkins, a Northern Italian (Ligurian) minority in south-western Sardinia. Gen. Res. Crop. Evol. 2008, 55, 911–924.
 11. Şenkardeş, I.; Tuzlacı, E. Some ethnobotanical notes from Gündoğmuş District (Antalya/Turkey). Clin. Exp. Health Sci. 2014, 4, 63–75.
 12. Sisay, M.; Gashaw, T. Ethnobotanical, ethnopharmacological, and phytochemical studies of *Myrtus communis* Linn: A popular herb in Unani system of medicine. J. Evid.-Based Complem. Altern. Med. 2017, 22, 1035–1043.
 13. Asllani, U. Chemical composition of albanian myrtle oil (*Myrtus communis* L.). J. Essent. Oil Res. 2000, 12, 140–142.
 14. Berka-Zougali, B.; Ferhat, M.A.; Hassani, A.; Chemat, F.; Allaf, K.S. Comparative study of essential oils extracted from Algerian *Myrtus communis* L. leaves using microwaves and hydrodistillation. Int. J. Mol. Sci. 2012, 13, 4673–4695.
 15. Bouzabata, A.; Casanova, J.; Bighelli, A.; Cavaleiro, C.; Salgueiro, L.; Tomi, F. The genus *Myrtus* L. in Algeria: Composition and biologicala of essential oils from *M. communis* and *M. nivellei*: A review. Chem. Biodivers. 2016, 13, 672–680.
 16. Tuberoso, C.I.; Barra, A.; Angioni, A.; Sarritzu, E.; Pirisi, F.M. Chemical composition of volatiles in Sardinian myrtle (*Myrtus communis* L.) alcoholic extracts and essential oils. J. Agric. Food Chem. 2006, 54, 1420–1426.
 17. Usai, M.; Mulas, M.; Marchetti, M. Chemical composition of essential oils of leaves and flowers from five cultivars of myrtle (*Myrtus communis* L.). J. Essent. Oil Res. 2015, 27, 465–476.
 18. Taamalli, A.; Iswaldi, I.; Arráez-Román, D.; Segura-Carretero, A.; Fernández-Gutiérrez, A.; Zarrouk, M. (2013). UPLC–QTOF/MS for a rapid characterisation of phenolic compounds from Leaves of *Myrtus communis* L. Phytochem. Anal. 25: 89–96.
 19. Yoshimura, M.; Amakura, Y.; Tokuhara, M.; Yoshida, T. (2008). Polyphenolic compounds isolated from the leaves of *Myrtus communis*. J. Nat. Med., 62: 366–368.
 20. Wannes, W.A.; Mhamdi, B.; Sriti, J.; Jemia, M.B.; Ouchikh, O.; Hamdaoui, G.; Kchouk, M.E.; Marzouk, B. (2010). Antioxidant activities of the essential oils and methanol extracts from myrtle (*Myrtus communis* var. *italica* L.) leaf, stem and flower. Food Chem. Toxicol. 48: 1362–1370.
 21. Harassi, Y.; Tilaoui, M.; Idir, A.; Frédéric, J.; Baudino, S.; Ajouaoi, S.; Mouse, H.A.; Zyad, A. (2019). Phytochemical analysis, cytotoxic and antioxidant activities of *Myrtus communis* essential oil from Morocco. J. Complement. Integr. Med., 16: 20180100.

22. Alipour, G.; Dashti, S.; Hosseinzadeh, H. (2014). Review of pharmacological effects of *Myrtus communis* L. and its active constituents. *Phytother. Res.* 28, 1125–1136.
23. Cannas, S.; Molicotti, P.; Ruggeri, M.; Cubeddu, M.; Sanguinetti, M.; Marongiu, B.; Zanetti, S. (2013). Antimycotic activity of *Myrtus communis* L. towards *Candida* spp. from clinical isolates. *J. Infect. Dev. Ctries.*, 7: 295–298.
24. Barac, A.; Donadu, M.; Usai, D.; Spiric, V.T.; Mazzarello, V.; Zanetti, S.; Aleksic, E.; Stevanovic, G.; Nikolic, N.; Rubino, S. (2018). Antifungal activity of *Myrtus communis* against *Malassezia* spp. isolated from the skin of patients with pityriasis versicolor. *Infection.* 46: 253–257.
25. Cannas, S.; Molicotti, P.; Usai, D.; Maxia, A.; Zanetti, S. (2014). Antifungal, anti-biofilm and adhesion activity of the essential oil of *Myrtus communis* L. against *Candida* species. *Nat. Prod. Res.* 28, 2173–2177.
26. Nabavizadeh, M.; Abbaszadegan, A.; Gholami, A.; Sheikhi, R.; Shokouhi, M.; Shams, M.S.; Ghasemi, Y. (2014). Chemical constituent and antimicrobial effect of essential oil from *Myrtus communis* leaves on microorganisms involved in persistent endodontic infection compared to two common endodontic irrigants: An in vitro study. *J. Conserv. Dent.*, 17: 449.
27. Mahboubi, M.; Bidgoli, F.G. (2010). In vitro synergistic efficacy of combination of amphotericin B with *Myrtus communis* essential oil against clinical isolates of *Candida albicans*. *Phytomedicine*, 17: 771–774.
28. Mahmoudvand, H.; Ezzatkhan, F.; Sharififar, F.; Sharifi, I.; Dezaki, E.S. (2015). Antileishmanial and cytotoxic effects of essential oil and methanolic extract of *Myrtus communis* L. *Korean J. Parasitol.* 53: 21.
29. Jabri, M.A.; Marzouki, L.; Sebai, H. Ethnobotanical, phytochemical and therapeutic effects of *Myrtus communis* L. berries seeds on gastrointestinal tract diseases: A review. *Arch. Physiol. Biochem.* 2018, 124, 390–396.
30. Mulas M.; Melis R.A.M. (2008). Influence of growing area, year, season, and cultivar on the composition of myrtle leaves and infusions *HortScience*, 43 (2): 549–553.
31. Anwar S.; Crouch R.A.; Awadh A.; Al-Fatimi M.A.; Setzer W.N.; Wessjohann L. (2017). Hierarchical cluster analysis and chemical characterisation of *Myrtus communis* L. essential oil from Yemen region and its antimicrobial, antioxidant and anti-colorectal adenocarcinoma properties. *J. Nat. Product Res.* 31: 2158–2163.
32. Hayder N, Abdelwaheda A, Kilania S, Ben Ammar R, Mahmoud A and Ghedirab K. (2004). Anti-genotoxic and free-radical scavenging activities of extracts from (Tunisian) *Myrtus communis*, *Mutat Res*, 564, 89–95.
33. Khurshid C A, Karrar M, Osamah I A, Rima O, Mostafizur R. and Coen R. (2022). Assessment of Potentially Toxic Elements in the Urban Soil and Plants of Kirkuk City in Iraq. *Sustainability.* 14(9), 5655.
34. Rastogi R P and Mehrotra B N, (1991). *Compendium of Indian Medicinal Plants (1970–1979)*, Central Drug Research Institute Lucknow, Vol. 2, Publications and Information Directorate, CSIR, New Delhi, p. 478.
35. Serce S, Ercisli S, Sengul M, Gunduz K and Orhan E. (2010). Antioxidant activities and fatty acid composition of wild grown myrtle (*Myrtus communis* L.) fruits, *Phcog Mag*, 6, 9–12.

36. Giacomo M, (1983). Gas chromatographic-mass spectrometric investigation of the volatile components of myrtle berries (*Myrtus communis* L.), J Chromatogr, 264, 304-311.
37. Jerkovic I, Radionic A and Borcic I, (2002). Comparative study of leaf, fruit and flower essential oils of Croatian *Myrtus communis* Linn. during a one year vegetative cycle, J. Essent Oil Res, 14 (4), 266-270.
38. Masoudi M, Miraj S, Rafieian-Kopaei M. (2016). Comparison of the Effects of *Myrtus Communis* L, *Berberis Vulgaris* and Metronidazole Vaginal Gel alone for the Treatment of Bacterial Vaginosis. J Clin Diagn Res. 10(3):QC04–7.
39. Ebrahimabadi EH, Ghoreishi SM, Masoum S, Ebrahimabadi AHJ (2016). Combination of GC/FID/Mass spectrometry fingerprints and multivariate calibration techniques for recognition of antimicrobial constituents of *Myrtus communis* L. essential oil. G. Chromatogr B Analyt Technol Biomed Life Sci. 1008:50-7.
40. Nabavizadeh M, Abbaszadegan A, Gholami A, Sheikhiani R, Shokouhi M, Shams MS, et al. (2014). Chemical constituent and antimicrobial effect of essential oil from *Myrtus communis* leaves on microorganisms involved in persistent endodontic infection compared to two common endodontic irrigants: An in vitro study. J Conserv Dent. 17(5):449-53.
41. Mohammadi R, Esfahani S H M, Shadzi S and Moattar F. (2008). Antifungal activity of *Myrtus communis* L. essential oil against clinical isolates of *Aspergillus*, J Isfahan Medical School, 26(89): 105-111.
42. Kordali S, Usanmaz A, Cakir A, Komaki A, Ercisli S. (2016). Antifungal and herbicidal effects of fruit essential oils of four *Myrtus communis* genotypes. Chem Biodivers 13 (1): 77-84.
43. Serce S., Ercisli S., Sengul M., Gunduz K., Orhan E. (2010). Antioxidant activities and fatty acid composition of wild grown myrtle (*Myrtus communis* L.) fruits. Pharmacogn. Mag. 6: 1-7.
44. Polat B., Oba S., Karaman K., Arici M., Sagdic O. (2014). Comparison of different solvent types for determination biological activities of myrtle berries collected from Turkey. Qual. Assur. Saf. Crops. 6: 221–227.
45. Messaoud C., Boussaid M. (2011). *Myrtus communis* berry color morphs: A comparative analysis of essential oils, fatty acids, phenolic compounds, and antioxidant activities. Chem. Biodivers. 8:300–310.
46. Tuberoso C.I.G., Rosa A., Bifulco E., Melis M.P., Atzeri A., Pirisi F.M., Dessì M.A. (2010). Chemical composition and antioxidant activities of *Myrtus communis* L. berries extracts. Food Chem. 123:1242–1251.
47. Al-zohyri A M, Al-Jeboory A A and Jawad A L M. (1985). Cardiovascular and antimicrobial effect of *Myrtus communis*, Indian J Pharmacol, 17(4), 233-235.
48. Kumar R, Phaneendra P, Bodhanapu S, Rahiman F, Niay M & Tamizmani T. (2011). Antioxidant and hepato-protective activity of the aqueous extract of *Myrtus communis* (myrtle) leaves, Pharmacology, 1: 1083-1090.
49. Zaidi SFH, Yamada K, Kadowaki M, Usmanghani K, Sugiyama T. (2009). Bactericidal activity of medicinal plants, employed for the treatment of gastrointestinal ailments, against *Helicobacter pylori*. J Ethnopharmacol 121: 286–291.

50. Sumbul S, Ahmad MA, Asif M, Akhtar M, Saud I. (2012). Physicochemical and phytochemical standardization of berries of *Myrtus communis* Linn. *J Pharm Bioall Sci* 4: 322–326.
51. Hosseinzadeh H, Khoshdel M, Ghorbani M. (2011). Antinociceptive, Anti-inflammatory Effects and Acute Toxicity of Aqueous and Ethanolic Extracts of *Myrtus communis* L. Aerial Parts in Mice. *J Acupunct Meridian Stud* 4: 242–247.
52. Sumbul S, Ahmad MA, Asif M, Saud I, Akhtar M. (2010). Evaluation of *Myrtus communis* Linn. berries (common myrtle) in experimental ulcer models in rats. *Hum Exp Toxicol* 29: 935–944.
53. Feisst C, Franke L, Appendino G and Werz O. (2005). Identification of molecular targets of the oligomeric nonprenylated acylphloroglucinols from *Myrtus communis* and their implication as anti-inflammatory compounds, *J Pharmacol Exp Ther*, 315(1): 389-396.
54. Ayvaz A, Sagdic O, Karaborklu S and Ozturk I. (2008). Insecticidal activity of the essential oils from different plants against three stored-product insects, *J Insect Sci*, 10(21), 1-13.
55. Gautheir R, Agoumi A and Gourai M. (1988). Activity of the extracts of *Myrtus communis* against *Pediculus humanis capitis*, *Plant Med Phytother*, 23(2), 25-108.
56. Mimica-Dukić N, Bugarin D, Grbović S, et al. (2010). Essential oil of *Myrtus communis* L. as a potential antioxidant and antimutagenic agents. *Molecules* 15: 2759–2770.

